

# Continuous Random Variables: Conditioning, Expectation and Independence



Berlin Chen  
Department of Computer Science & Information Engineering  
National Taiwan Normal University



## Reference:

- D. P. Bertsekas, J. N. Tsitsiklis, *Introduction to Probability*, Sections 3.4-3.5

# Conditioning PDF Given an Event (1/3)

- The conditional PDF of a continuous random variable  $X$ , given an event  $A$ 
  - If  $A$  cannot be described in terms of  $X$ , the conditional PDF is defined as a nonnegative function  $f_{X|A}(x)$  satisfying

$$\mathbf{P}(X \in B | A) = \int_B f_{X|A}(x) dx$$

- Normalization property

$$\int_{-\infty}^{\infty} f_{X|A}(x) dx = 1$$

## Conditioning PDF Given an Event (2/3)


- If  $A$  can be described in terms of  $X$  ( $A$  is a subset of the real line with  $\mathbf{P}(X \in A) > 0$ ), the conditional PDF is defined as a nonnegative function  $f_{X|A}(x)$  satisfying

$$f_{X|A}(x) = \begin{cases} \frac{f_X(x)}{\mathbf{P}(X \in A)}, & \text{if } x \in A \\ 0, & \text{otherwise} \end{cases}$$

and for any subset  $B$

$$\begin{aligned} \mathbf{P}(X \in B | X \in A) &= \frac{\mathbf{P}(X \in B, X \in A)}{\mathbf{P}(X \in A)} \\ &= \frac{\int_{A \cap B} f_X(x) dx}{\mathbf{P}(X \in A)} \\ &= \int_B f_{X|A}(x) dx \end{aligned}$$

- Normalization Property  $\int_{-\infty}^{\infty} f_{X|A}(x) dx = \int_A f_{X|A}(x) dx = 1$



$f_{X|A}$  remains the same shape as  $f_X$  except that it is scaled along the vertical axis

## Conditioning PDF Given an Event (3/3)

- If  $A_1, A_2, \dots, A_n$  are disjoint events with  $\mathbf{P}(A_i) > 0$  for each  $i$ , that form a partition of the sample space, then

$$f_X(x) = \sum_{i=1}^n \mathbf{P}(A_i) f_{X|A_i}(x)$$

- Verification of the above total probability theorem

$$\mathbf{P}(X \leq x) = \sum_{i=1}^n \mathbf{P}(A_i) \mathbf{P}(X \leq x | A_i)$$

$$\Rightarrow \int_{-\infty}^x f_X(t) dt = \sum_{i=1}^n \mathbf{P}(A_i) \int_{-\infty}^x f_{X|A_i}(t) dt$$

Taking the derivative of both sides with respective to  $x$

$$\Rightarrow f_X(x) = \sum_{i=1}^n \mathbf{P}(A_i) f_{X|A_i}(x)$$

# Conditional Expectation Given an Event

- The conditional expectation of a continuous random variable  $X$ , given an event  $A$  ( $P(A) > 0$ ), is defined by

$$E[X|A] = \int_{-\infty}^{\infty} xf_{X|A}(x)dx$$

- The conditional expectation of a function  $g(X)$  also has the form

$$E[g(X)|A] = \int_{-\infty}^{\infty} g(x)f_{X|A}(x)dx$$

- Total Expectation Theorem

$$E[g(X)] = \sum_{i=1}^n P(A_i)E[g(X)|A_i]$$


- Where  $A_1, A_2, \dots, A_n$  are disjoint events with  $P(A_i) > 0$  for each  $i$ , that form a partition of the sample space

# An Illustrative Example

- **Example 3.10. Mean and Variance of a Piecewise Constant PDF.**

Suppose that the random variable  $X$  has the piecewise constant PDF

$$f_X(x) = \begin{cases} 1/3, & \text{if } 0 \leq x \leq 1, \\ 2/3, & \text{if } 1 \leq x \leq 2, \\ 0, & \text{otherwise.} \end{cases}$$



Define event  $A_1 = \{X \text{ lies in the first interval } [0,1]\}$

event  $A_2 = \{X \text{ lies in the second interval } [1,2]\}$

$$\Rightarrow \mathbf{P}(A_1) = \int_0^1 1/3 dx = 1/3, \quad \mathbf{P}(A_2) = \int_1^2 2/3 dx = 2/3$$

Recall that the mean and second moment of a uniform random variable over an interval  $[a, b]$  is  $(a + b)/2$  and  $(a^2 + ab + b^2)/3$

$$\Rightarrow \mathbf{E}[X|A_1] = 1/2, \quad \mathbf{E}[X^2|A_1] = 1/3$$

$$\mathbf{E}[X|A_2] = 3/2, \quad \mathbf{E}[X^2|A_2] = 7/3$$

$$\begin{aligned} \Rightarrow \mathbf{E}[X] &= \mathbf{P}(A_1)\mathbf{E}[X|A_1] + \mathbf{P}(A_2)\mathbf{E}[X|A_2] \\ &= 1/3 \cdot 1/2 + 2/3 \cdot 3/2 = 7/6 \end{aligned}$$

$$\begin{aligned} \mathbf{E}[X^2] &= \mathbf{P}(A_1)\mathbf{E}[X^2|A_1] + \mathbf{P}(A_2)\mathbf{E}[X^2|A_2] \\ &= 1/3 \cdot 1/3 + 2/3 \cdot 7/3 = 15/9 \end{aligned}$$

$$\therefore \text{var}(X) = 15/9 - (7/6)^2 = 11/36$$

# Multiple Continuous Random Variables

- Two continuous random variables  $X$  and  $Y$  associated with a common experiment are **jointly continuous** and can be described in terms of a **joint PDF**  $f_{X,Y}$  satisfying

$$\mathbf{P}((X, Y) \in B) = \iint\limits_{(x, y) \in B} f_{X,Y}(x, y) dx dy$$

- $f_{X,Y}$  is a nonnegative function
  - Normalization Probability  $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx dy = 1$
- Similarly,  $f_{X,Y}(a, c)$  can be viewed as the “probability per unit area” in the vicinity of  $(a, c)$

$$\begin{aligned}\mathbf{P}(a \leq X \leq a + \delta, c \leq Y \leq c + \delta) \\ = \int_a^{a+\delta} \int_c^{c+\delta} f_{X,Y}(x, y) dx dy = f_{X,Y}(a, c) \cdot \delta^2\end{aligned}$$

- Where  $\delta$  is a small positive number

# An Illustrative Example

- **Example 3.13. Two-Dimensional Uniform PDF.** We are told that the joint PDF of the random variables  $X$  and  $Y$  is a constant  $c$  on an area  $S$  and is zero outside. Find the value of  $c$  and the marginal PDFs of  $X$  and  $Y$ .

The corresponding uniform joint PDF on an area  $S$  is defined to be (cf. Example 3.12)

$$f_{X,Y}(x,y) = \begin{cases} \frac{1}{\text{Size of area } S}, & \text{if } (x,y) \in S \\ 0, & \text{otherwise} \end{cases}$$

$$\Rightarrow f_{X,Y}(x,y) = \frac{1}{4} \text{ for } (x,y) \in S$$

for  $1 \leq x \leq 2$

$$\begin{aligned} \Rightarrow f_X(x) &= \int_1^4 f_{X,Y}(x,y) dy \\ &= \int_1^4 \frac{1}{4} dy = \frac{3}{4} \end{aligned}$$

for  $2 \leq x \leq 3$


$$\begin{aligned} \Rightarrow f_X(x) &= \int_2^3 f_{X,Y}(x,y) dy \\ &= \int_2^3 \frac{1}{4} dy = \frac{1}{4} \end{aligned}$$

for  $1 \leq y \leq 2$

$$\begin{aligned} \Rightarrow f_Y(y) &= \int_1^2 f_{X,Y}(x,y) dx \\ &= \int_1^2 \frac{1}{4} dx = \frac{1}{4} \end{aligned}$$

for  $2 \leq y \leq 3$

$$\begin{aligned} \Rightarrow f_Y(y) &= \int_1^3 f_{X,Y}(x,y) dx \\ &= \int_1^3 \frac{1}{4} dx = \frac{1}{2} \end{aligned}$$



for  $3 \leq y \leq 4$

$$\begin{aligned} \Rightarrow f_Y(y) &= \int_1^2 f_{X,Y}(x,y) dx \\ &= \int_1^2 \frac{1}{4} dx = \frac{1}{4} \end{aligned}$$

# Conditioning one Random Variable on Another

- Two continuous random variables  $X$  and  $Y$  have a joint PDF. For any  $y$  with  $f_Y(y) > 0$ , the conditional PDF of  $X$  given that  $Y = y$  is defined by

$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)}$$


- Normalization Property  $\int_{-\infty}^{\infty} f_{X|Y}(x|y)dx = 1$
- The marginal, joint and conditional PDFs are related to each other by the following formulas

$$f_{X,Y}(x,y) = f_Y(y)f_{X|Y}(x|y),$$

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y)dy. \quad \text{marginalization}$$

## Illustrative Examples (1/2)

- Notice that the conditional PDF  $f_{X|Y}(x|y)$  has the same shape as the joint PDF  $f_{X,Y}(x,y)$ , because the normalizing factor  $f_Y(y)$  does not depend on  $x$



$$f_{X|Y}(x|3.5) = \frac{f_{X,Y}(x, 3.5)}{f_Y(3.5)} = \frac{1/4}{1/4} = 1$$

$$f_{X|Y}(x|2.5) = \frac{f_{X,Y}(x, 2.5)}{f_Y(2.5)} = \frac{1/4}{1/2} = 1/2$$

$$f_{X|Y}(x|1.5) = \frac{f_{X,Y}(x, 1.5)}{f_Y(1.5)} = \frac{1/4}{1/4} = 1$$

cf. example 3.13

**Figure 3.17:** Visualization of the conditional PDF  $f_{X|Y}(x|y)$ .

Let  $X, Y$  have a joint PDF which is uniform on the set  $S$ . For each fixed  $y$ , we consider the joint PDF along the slice  $Y = y$  and normalize it so that it integrates to 1

## Illustrative Examples (2/2)


- **Example 3.15. Circular Uniform PDF.** Ben throws a dart at a circular target of radius  $r$ . We assume that he always hits the target, and that all points of impact  $(x, y)$  are equally likely, so that the joint PDF  $f_{X,Y}(x, y)$  of the random variables  $x$  and  $y$  is uniform
  - What is the marginal PDF  $f_Y(y)$

$$f_{X,Y}(x, y) = \begin{cases} \frac{1}{\text{area of the circle}}, & \text{if } (x, y) \text{ is in the circle} \\ 0, & \text{otherwise} \end{cases}$$

$$= \begin{cases} \frac{1}{\pi r^2}, & x^2 + y^2 \leq r^2 \\ 0, & \text{otherwise} \end{cases}$$

$$\begin{aligned} f_Y(y) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx = \int_{x^2 + y^2 \leq r^2} \frac{1}{\pi r^2} dx \\ &= \frac{1}{\pi r^2} \int_{x^2 + y^2 \leq r^2} 1 dx = \frac{1}{\pi r^2} \int_{-\sqrt{r^2 - y^2}}^{\sqrt{r^2 - y^2}} 1 dx \\ &= \frac{2}{\pi r^2} \sqrt{r^2 - y^2}, \text{ if } |y| \leq r \end{aligned}$$

(Notice here that PDF  $f_Y(y)$  is not uniform)



$$\begin{aligned} f_{X|Y}(x|y) &= \frac{f_{X,Y}(x,y)}{f_Y(y)} \\ &= \frac{\frac{1}{\pi r^2}}{\frac{2}{\pi r^2} \sqrt{r^2 - y^2}} \\ &= \frac{1}{2 \sqrt{r^2 - y^2}}, \quad \text{if } x^2 + y^2 \leq r^2 \end{aligned}$$

For each value  $y$ ,  $f_{X|Y}(x|y)$  is uniform

# Expectation of a Function of Random Variables

- If  $X$  and  $Y$  are jointly continuous random variables, and  $g$  is some function, then  $Z = g(X, Y)$  is also a random variable (can be continuous or discrete)
  - The expectation of  $Z$  can be calculated by

$$E[Z] = E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy$$

- If  $Z$  is a linear function of  $X$  and  $Y$ , e.g.,  $Z = aX + bY$ , then

$$E[Z] = E[aX + bY] = aE[X] + bE[Y]$$

- Where  $a$  and  $b$  are scalars

# Conditional Expectation

- The properties of **unconditional expectation** carry through, with the obvious modifications, to **conditional expectation**

$$E[X|Y=y] = \int_{-\infty}^{\infty} xf_{X|Y}(x|y)dx$$

$$E[g(X)|Y=y] = \int_{-\infty}^{\infty} g(x)f_{X|Y}(x|y)dx$$

$$E[g(X,Y)|Y=y] = \int_{-\infty}^{\infty} g(x,y)f_{X|Y}(x|y)dx$$

# Total Probability/Expectation Theorems

- **Total Probability Theorem**

- For any event  $A$  and a continuous random variable  $Y$

$$\mathbf{P}(A) = \int_{-\infty}^{\infty} \mathbf{P}(A|Y = y) f_Y(y) dy$$

- **Total Expectation Theorem**

- For any continuous random variables  $X$  and  $Y$

$$\mathbf{E}[X] = \int_{-\infty}^{\infty} \mathbf{E}[X|Y = y] f_Y(y) dy$$

$$\mathbf{E}[g(X)] = \int_{-\infty}^{\infty} \mathbf{E}[g(X)|Y = y] f_Y(y) dy$$

$$\mathbf{E}[g(X, Y)] = \int_{-\infty}^{\infty} \mathbf{E}[g(X, Y)|Y = y] f_Y(y) dy$$

# Independence

- Two continuous random variables  $X$  and  $Y$  are independent if

$$f_{X,Y}(x,y) = f_X(x)f_Y(y), \text{ for all } x,y$$

– Or

$$f_{X|Y}(x|y) = f_X(x), \text{ for all } x \text{ and all } y \text{ with } f_Y(y) > 0$$

– Or

$$f_{Y|X}(y|x) = f_Y(y), \text{ for all } y \text{ and all } x \text{ with } f_X(x) > 0$$

## More Factors about Independence (1/2)

- If two continuous random variables  $X$  and  $Y$  are independent, then
  - Any two events of the forms  $\{X \in A\}$  and  $\{Y \in B\}$  are independent

$$\begin{aligned}\mathbf{P}(X \in A, Y \in B) &= \int_{x \in A} \int_{y \in B} f_{X,Y}(x, y) dy dx \\ &= \int_{x \in A} \int_{y \in B} f_X(x) f_Y(y) dy dx \\ &= \left[ \int_{x \in A} f_X(x) dx \right] \left[ \int_{y \in B} f_Y(y) dy \right] \\ &= \mathbf{P}(X \in A) \mathbf{P}(Y \in B)\end{aligned}$$

## More Factors about Independence (2/2)

- If two continuous random variables  $X$  and  $Y$  are independent, then
  - $\mathbf{E}[XY] = \mathbf{E}[X]\mathbf{E}[Y]$
  - $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y)$
  - The random variables  $g(X)$  and  $h(Y)$  are independent for any functions  $g$  and  $h$ 
    - Therefore,

$$\mathbf{E}[g(X)h(Y)] = \mathbf{E}[g(X)]\mathbf{E}[h(Y)]$$

## Joint CDFs

- If  $X$  and  $Y$  are two (either continuous or discrete) random variables, their joint cumulative distribution function (CDF) is defined by

$$F_{X,Y}(x, y) = \mathbf{P}(X \leq x, Y \leq y)$$

- If  $X$  and  $Y$  further have a joint PDF  $f_{X,Y}$ , then

$$F_{X,Y}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{X,Y}(s, t) ds dt$$

And


$$f_{X,Y}(x, y) = \frac{\partial^2 F_{X,Y}(x, y)}{\partial x \partial y}$$

If  $F_{X,Y}$  can be differentiated at the point  $(x, y)$

## An Illustrative Example

- **Example 3.20.** Verify that if  $X$  and  $Y$  are described by a uniform PDF on the unit square, then the joint CDF is given by

$$F_{X,Y}(x,y) = \mathbf{P}(X \leq x, Y \leq y) = xy, \quad \text{for } 0 \leq x, y \leq 1$$



$$\frac{\partial^2 F_{X,Y}(x,y)}{\partial x \partial y} = 1 = f_{X,Y}(x,y), \quad \text{for all } (x,y) \text{ in the unit square}$$

## Recall: the Discrete Bayes' Rule


- Let  $A_1, A_2, \dots, A_n$  be disjoint events that form a partition of the sample space, and assume that  $P(A_i) \geq 0$  for all  $i$ . Then, for any event  $B$  such that  $P(B) > 0$  we have

$$\begin{aligned} P(A_i|B) &= \frac{P(A_i)P(B|A_i)}{P(B)} \\ &= \frac{P(A_i)P(B|A_i)}{\sum_{k=1}^n P(A_k)P(B|A_k)} \\ &= \frac{P(A_i)P(B|A_i)}{P(A_1)P(B|A_1) + \dots + P(A_n)P(B|A_n)} \end{aligned}$$


Multiplication rule  
Total probability theorem

# Inference and the Continuous Bayes' Rule (1/2)

- As we have a model of an underlying but unobserved phenomenon, represented by a random variable  $X$  with PDF  $f_X$ , and we make a noisy measurement  $Y$ , which is modeled in terms of a conditional PDF  $f_{Y|X}$ . Once the experimental value of  $Y$  is measured, what information does this provide on the unknown value of  $X$ ?



$$f_{X|Y}(x|y) = \frac{f_{X,Y}(x,y)}{f_Y(y)} = \frac{f_X(x)f_{Y|X}(y|x)}{\int_{-\infty}^{\infty} f_X(t)f_{Y|X}(y|t)dt}$$

## Inference and the Continuous Bayes' Rule (2/2)

- If the unobserved phenomenon is inherently discrete
  - Let  $N$  is a discrete random variable of the form  $\{N = n\}$  that represents the different discrete probabilities for the unobserved phenomenon of interest, and  $p_N$  be the PMF of  $N$

$$\begin{aligned}\mathbf{P}(N = n | Y = y) &\approx \mathbf{P}(N = n | y \leq Y \leq y + \delta) \\ &= \frac{\mathbf{P}(N = n)\mathbf{P}(y \leq Y \leq y + \delta | N = n)}{\mathbf{P}(y \leq Y \leq y + \delta)} \\ &\approx \frac{p_N(n)f_{Y|N}(y|n)\delta}{f_Y(y)\delta} \quad \text{Total probability theorem} \\ &= \frac{p_N(n)f_{Y|N}(y|n)}{\sum_i p_N(i)f_{Y|N}(y|n)}\end{aligned}$$

## Illustrative Examples (1/2)

- **Example 3.18.** A lightbulb produced by the General Illumination Company is known to have an exponentially distributed lifetime  $Y$ . However, the company has been experiencing quality control problems. On any given day, the parameter  $\Lambda = \lambda$  of the PDF of  $Y$  is actually a random variable, uniformly distributed in the interval  $[1, 3/2]$ .
  - If we test a lightbulb and record its lifetime ( $Y = y$ ), what can we say about the underlying parameter  $\lambda$ ?

$$f_{Y|\Lambda}(y|\lambda) = \lambda e^{-\lambda y}, \quad y \geq 0, \lambda > 0$$

Conditioned on  $\Lambda = \lambda$ ,  $Y$  has a exponential distribution with parameter  $\lambda$

$$f_\Lambda(\lambda) = \begin{cases} 2, & \text{for } 1 \leq \lambda \leq 3/2 \\ 0, & \text{otherwise} \end{cases}$$

$$f_{\Lambda|Y}(\lambda|y) = \frac{f_\Lambda(\lambda)f_{Y|\Lambda}(y|\lambda)}{\int_1^{3/2} f_\Lambda(t)f_{Y|\Lambda}(y|t)dt} = \frac{2\lambda e^{-\lambda y}}{\int_1^{3/2} 2te^{-ty} dt}, \quad \text{for } 1 \leq \lambda \leq 3/2$$

## Illustrative Examples (2/2)

- **Example 3.19. Signal Detection.** A binary signal  $S$  is transmitted, and we are given that  $\mathbf{P}(S = 1) = p$  and  $\mathbf{P}(S = -1) = 1 - p$ .
  - The received signal is  $Y = S + N$ , where  $N$  normal noise with zero mean and unit variance, independent of  $S$ .
  - What is the probability that  $S = 1$ , as a function of the observed value  $y$  of  $Y$ ?

$$f_{Y|S}(y|s) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(y-s)^2/2}, \text{ for } s = 1 \text{ and } -1, \text{ and } -\infty \leq y \leq \infty$$

Conditioned on  $S = s$ ,  $Y$  has a normal distribution with mean  $s$  and unit variance

$$\begin{aligned} \mathbf{P}(S = 1|Y = y) &= \frac{p_S(1)f_{Y|S}(y|1)}{f_Y(y)} = \frac{p_S(1)f_{Y|S}(y|1)}{p_S(1)f_{Y|S}(y|1) + p_S(-1)f_{Y|S}(y|-1)} \\ &= \frac{p \frac{1}{\sqrt{2\pi}} e^{-(y-1)^2/2}}{p \frac{1}{\sqrt{2\pi}} e^{-(y-1)^2/2} + (1-p) \frac{1}{\sqrt{2\pi}} e^{-(y+1)^2/2}} \\ &= \frac{e^{-(y^2+1)/2} \cdot pe^y}{e^{-(y^2+1)/2} \cdot pe^y + e^{-(y^2+1)/2} \cdot (1-p)e^{-y}} = \frac{pe^y}{pe^y + (1-p)e^{-y}} \end{aligned}$$

# Recitation

- SECTION 3.4 Conditioning on an Event
  - Problems 14, 17, 18
- SECTION 3.5 Multiple Continuous Random Variables
  - Problems 19, 24, 25, 26, 28